

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : A61K 33/06, 31/20, 31/22, 31/52, 31/55		A1	(11) International Publication Number: WO 00/66133
			(43) International Publication Date: 9 November 2000 (09.11.00)
(21) International Application Number: PCT/US00/11681		(81) Designated States: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).	
(22) International Filing Date: 28 April 2000 (28.04.00)			
(30) Priority Data: 09/303,689 3 May 1999 (03.05.99) US			
(71) Applicant: KV PHARMACEUTICAL COMPANY [US/US]; 2503 South Hanley Road, St. Louis, MO 63144-2555 (US).			
(72) Inventors: HERMELIN, Marc, S.; 791 Bismark, Glendale, MO 63122 (US). LEVINSON, R., Saul; 17020 Kimwood Court, Chesterfield, MO 63005 (US).			
(74) Agent: NATH, Gary, M.; Nath & Associates PLLC, 6th floor, 1030 15th Street, N.W., Washington, DC 20005-1503 (US).			
Published <i>With international search report.</i>			
(54) Title: DIETARY SUPPLEMENT			
(57) Abstract <p>The present disclosure relates to novel nutritional compositions and methods for augmenting the possibility of conception while increasing nutritional stores to aid development of healthy embryos and child growth. The nutritional compositions are intended for use by both males and females planning to conceive a child.</p>			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

DIETARY SUPPLEMENT**BACKGROUND OF THE INVENTION****Field of the Invention**

The present invention is directed to novel
5 nutritional compositions, particularly nutritional
compositions for men and women planning to
conceive children, and methods of using said
compositions prior to and during pregnancy to
augment the possibility of conception occurring
10 and/or increase nutritional stores for aiding the
development of healthy embryos and child growth.

Description of the Related Art

Infertility is a serious problem in the
15 United States and throughout the world, in both
industrialized and unindustrialized nations. In
the United States alone, infertility affects an
estimated 20 million families (i.e., approximately
20% of all U.S. families). See *The Merck Manual*,
20 1768 (16th Ed. 1992). In about 40% of these cases,
the infertility is attributable to the male and in
about 40-50% of these cases, the infertility is
attributable to the female (note: the cause of
infertility in about 10-20% of the cases is
25 indeterminate). See *McGraw-Hill Encyclopedia of
Science and Technology*, 17:417 (6th Ed. 1987).

Evidence indicates that the general health of both males and females prior to conception has a direct impact upon the ability to conceive. See *Understanding Nutrition*, 479-480 (Whitney and Rolfes eds., 6th Ed., 1993). Further, studies of both men and women have shown that the underlying cause of infertility in a marked proportion of individuals may be attributed to a nutritional factor. See *Id.* In fact, it has been suggested that the inability to reproduce is one of the first signs of imperfect nutrition. See *Id.*

Infertility in men is primarily associated with low sperm count, decreased sperm motility, sperm agglutination, impotence and ejaculatory disorders. See *The Merck Manual*, 1768 (16th Ed. 1992). Animal studies suggest that dietary ascorbate (vitamin C) levels directly affect sperm quality and influence male fertility in scurvy-prone vertebrates. It is believed that high concentrations of ascorbic acid in semen play a key role in maintaining the genetic integrity of sperm cells by preventing oxidative damage to sperm DNA. See Dabrowski, "Ascorbic acid protects against male infertility in teleost fish", *Experientia*, 52(2):97-100 (1996). There is also

evidence that daily vitamin C therapy is useful in the treatment and/or mitigation of decreased sperm motility and agglutination. Gonzalez, "Sperm swim singly after vitamin C therapy", *JAMA*, 249(20):2747, 2751 (1983).

Various studies suggest that vitamin E is also effective in treating male infertility. For example, one study involving the oral dosing of vitamin E over a three month period resulted in a 50 percent increase in spermatozoal zona binding. See Kessopoulou, "A double-blind randomizing placebo cross-over controlled trial using the antioxidant vitamin E to treat reactive oxygen species associated male infertility", *Fertil Steril*, 64(4): 825-31 (1995). Another study found that treatment of male infertility patients with oral vitamin E significantly decreased malondialdehyde concentrations, high levels of which are indicia of decreased sperm motility. Suleiman, "Lipid peroxidation and human sperm motility: protective role of vitamin E", *J Androl*, 17(5):530-7 (1996); See also, Vezina, "Selenium-vitamin E supplementation in infertile men. Effects on semen parameters and micronutrient levels and distribution", *Bio Trace Elem Res*,

53(1-3):65-83 (1996).

Infertility in women is primarily associated with dysfunction of ovulation, abnormal fallopian tube function and low cervical mucus receptivity.

5 *The Merck Manual*, 1770-1772 (16th Ed. 1992).
Infertility in women has also been linked to abnormally low red cell magnesium levels, and such cases have been successfully treated with dietary supplementation of oral magnesium. See Howard,
10 "Red cell magnesium and glutathione peroxidase in infertile women-effects of oral supplementation with magnesium and selenium", *Magnes Res*, 7(1):49-57 (1994).

For women planning to conceive children, the
15 role of nutrition is not limited to infertility alone. A mother's body provides the environment in which development of the embryo and fetus occur. See *Understanding Nutrition*, 479-480 (Whitney and Rolfes eds., 6th ed., 1993).
20 Accordingly, a mother's nutritional status prior to conception directly impacts the development of the fetus and embryo and is therefore implicated in the risk of birth defects. See *Id.*

In particular, during the first 20-25 days of
25 pregnancy, the placenta is not yet formed and

fetal circulation is not yet established. Therefore, during this period the fetus is nourished via digested maternal uterine cells and the diffusion of blood exudates. See Schorah, "Importance of Adequate Folate Nutrition in Embryonic and Early Fetal Development," *Vitamins and Minerals in Pregnancy and Lactation*, 167-176 (Berger, ed., Vol. 16, 1988). Thus, it has been suggested that good nutrient supply is not only required in the very early stages of pregnancy, but also in the preconceptional period. See *Id.* It is believed that a good nutrient supply during the preconceptional period and first 20 to 25 days of pregnancy (i.e., the "histiotrophic nutritional phase") is necessary to provide optimal concentrations of essential micronutrients to the endometrium, into which the embryo will embed. See *Id.* Further, inadequate nutrition prior to and at the time of conception causes the placenta, the function of which is to nourish the developing fetus, to develop incorrectly. "Transplacental Nutrient Transfer and Intrauterine Growth Retardation," *Nutrition News* 50 (1992): 56-57.

Increased occurrences of birth defects have been linked to inadequate nutrition in women at

the time of conception. Cases of infants born with neural tube defect (NTD), i.e., spina bifida and anacephaly, have been documented in women with various nutritional deficiencies, primarily low blood folic acid and vitamin C concentrations. Smithells, "Vitamin deficiencies and neural tube defects", *Arch Dis Child* 51:944-50 (1976).

Various studies point to a correlation between certain vitamin and mineral deficiencies and the etiologies of specific disease states in humans. See, e.g., Diplock, "Antioxidant Nutrients and Disease Prevention: An Overview," *Am. J. Clin. Nutr.*, 53:189-193 (1991); *Documenta Giegy Scientific Tables*, 457-497 (Diem and Cemtuer eds., 7th ed., 1975). In particular, studies designed to test the causal relationship between specific micronutrient deficiencies and resulting birth defects elucidate a correlation between proper folic acid and vitamin C levels and the reduction in the recurrence of NTD in the instances where women have experienced at least one prior pregnancy resulting in a child with NTD. See Schorah, "Importance of adequate folate ntrition in embryonic and early fetal development", *Vitamins and Minerals in Pregnancy*

and *Lactation*, 167-176 (Berger, ed., Vol. 16, 1988).

Multi-vitamin and mineral supplements for treating specific medical conditions and as
5 general nutritional supplements to promote and maintain good health have been described in various references. In particular, compositions and methods for optimizing the general health of both men and women by supplementing the daily diet
10 with specific and multi-vitamin compositions are disclosed in the following references.

Jansen, U.S. Patent No. 4,945,083, describes multi-factor hematinic vitamin preparations which provide B₁₂ and folic acid in a one to one ratio in
15 fully effective daily dosage amounts.

Mehansho, U.S. Patent No. 4,994,283, describes nutritional mineral supplements comprised of iron compounds and calcium compounds in combination with citrates or tartrates,
20 ascorbates, and optionally, fructose, such that the tendency of calcium to inhibit the bioavailability of iron is reduced, and the conjoint bioavailability of these two important minerals is enhanced.

25 Briggs et al., U.S. Patent No. 4,752,479 also

describes a multi-vitamin and mineral dietary supplement composition for oral administration. The supplement contains one or more divalent dietary mineral components selected from the group consisting of bioavailable calcium and magnesium, optionally in the presence of one or more additional non-ferrous mineral and vitamin components adapted to be released in the upper gastrointestinal tract, and a bioavailable iron component, present in controlled release form and adapted to be slowly released lower in the gastrointestinal tract, and a method of preventing or treating iron deficiency using such compositions. The supplement could contain about 200 mg of bioavailable calcium and 50 mg of bioavailable magnesium.

Lawrence, U.S. Patent No. 4,931,441, describes a stabilized aqueous leucovorin calcium composition suitable for administration by injection. The solution can contain about 6.35 mg per mL of leucovorin calcium pentahydrate.

Multi-vitamin and mineral dietary vitamin supplements for pregnant and lactating women have also been described. Sultenfuss, U.S. Patent No. 5,514,382, discloses a daily vitamin and mineral

supplement for women which provides necessary nutrients to maintain present health as well as positively influence future health. The vitamin and mineral supplement comprises vitamin A, beta-carotene, niacin, riboflavin, pantothenic acid, pyridoxine, cyanocobalamin, biotin, para-aminobenzoic acid, inositol, choline, vitamin C, vitamin D, vitamin E, vitamin K, boron, calcium, chromium, copper, iodine, iron, magnesium, manganese, molybdenum, selenium, zinc, and bioflavonoid. Niacin (vitamin B₃) is present to facilitate the production of the majority of sex hormones by dilating blood vessels, lowering cholesterol and maintaining blood circulation.

Paradissis et al., U.S. Patent No. 5,494,678, disclose multi-vitamin and mineral supplements for pregnant women which are designed for administration during the first, second and third trimesters of pregnancy. The multi-vitamin and mineral supplements are comprised of a regime of pharmaceutically-acceptable calcium compounds including vitamin D, folic acid, vitamin B₁₂, vitamin B₆, and vitamin B₁. These prenatal supplements are specifically tailored to maximize fetal development and maternal health during each

trimester of pregnancy.

Guall, U.S. Patent No. 4,629,625, discloses both vitamin and/or mineral compositions containing taurine and the utilization of these compositions for human nutritional purposes either singularly or as a supplement to other vitamin and mineral regimes. According to the reference, the taurine-containing composition may be used in conjunction with prenatal vitamin supplements for use by pregnant and lactating mothers.

Uiterwaal et al., U.S. Patent No. 4,710,387, disclose nutritional supplement preparations intended for pregnant and breast-feeding women. The disclosed supplements are based upon milk constituents, including proteins, fats, carbohydrates, calcium, copper, zinc, iodine, iron, vitamin A, vitamin B₁, vitamin B₆, vitamin C, vitamin D₃, vitamin E, niacin and folic acid. The precise adjustments of the constituents of the preparation is determined by the current consumer.

Ellenbogen, U.S. Patent No. 4,431,634, discloses multimineral dietary supplement compositions of enhanced iron bioavailability containing magnesium and calcium for use in prenatal therapy specifically tailored to combat

iron-deficiency anemia.

The *Physicians' Desk Reference for Nonprescription Drugs* describes various vitamin and mineral supplements for use by women. For example, Stuart Prenatal® is a multi-vitamin and mineral supplement for use "before, during and after pregnancy." It provides vitamins equal to 100% or more of the RDA for pregnant and lactating women. See *Physicians' Desk Reference for Nonprescription Drugs*, (9th Ed., 1988) 712.

Materna® prenatal vitamin and mineral tablets are for use "before, during and after pregnancy" and is indicated to provide vitamin and mineral supplementation throughout pregnancy as well as the postnatal period for both lactating and nonlactating mothers. The reference states that Materna® is "useful for improving nutritional status prior to conception." *Physicians' Desk Reference*, (51st Ed., 1997) 1427.

PreCare® prenatal multi-vitamin and mineral supplement film coated caplets are indicated to provide vitamin and mineral supplementation throughout pregnancy and during postnatal period-both for lactating and nonlactating mothers. The reference discloses that PreCare® is

useful for improving nutritional status prior to conception." *Physicians' Desk Reference*, (51st Ed., 1997) 2753.

5 Methods and compositions for treating infertility have also been described in various references. Madsen, U.S. Patent No. 5,389,657, discloses methods of treating infertility by administering therapeutically effective amounts of the glutathione stimulator L-2-oxothiazolidine-4-
10 carboxylate or esters thereof to female mammals having fertility problems. According to the reference, the formulation could be used as part of a complete nutritional formulation for satisfying nutritional requirements.

15 The compositions and methods discussed above are deficient in various respects. First, the compositions are not specifically formulated to address the problem of infertility. Even the above discussed references which recognize the
20 correlation between nutritional status and infertility do not specifically disclose any specific nutritional formulations for treating infertility and do not offer any guidance with regard to formulating specific nutritional
25 compositions for treating infertility. Thus,

these references are inadequate with regard to improving the possibility of conception.

Secondly, previously disclosed compositions only provide nutritional components in levels indicated for maintaining general health and are not need-specific formulations which are designed to address the distinct nutritional needs of men and women planning to conceive. Further, even previously disclosed need-based multi-vitamin and mineral supplements are limited to addressing the general nutritional needs of pregnant women, lactating women or nonpregnant, or non-lactating women. Thus, the specific nutritional needs of men and women during the period of time just prior to conception are not addressed by conventional nutritional supplementation.

Thirdly, conventional ovulatory inducing agents or other infertility agents do not utilize vitamins and minerals as active components and are thus not effective in addressing nutritional needs of men and women planning to conceive, or in reducing the risks of birth defects. Further, these agents may increase the risk of birth defects or have other undesirable side effects.

Therefore, there remains a need for specific

nutritional formulations which augment the possibility of conception occurring and reduce the risk of birth defects, as well as support general health. Moreover, there is a particular need for formulations which simultaneously augment the possibility of conception and reduce the risk of birth defects to provide a higher degree of patient compliance and minimize the cost to the patients. Additionally, it is desirable to have formulations which specifically address the differing needs of males and females during the period of time prior to conception.

There is also a need for multi-vitamin and mineral supplements which provide a regimen specifically designed to meet the nutritional requirements of males and females planning to conceive by providing the dosages of vitamins and minerals necessary to avoid vitamin or mineral deficiencies, and in particular those deficiencies associated with male and female infertility. It is always desirable to have formulations which minimize the necessity for medications. It is also particularly desirable to have available formulations for addressing infertility which are suitable for men and women seeking to limit their

use of medications. Thus, there is a general overall need for a fundamentally new, safe, effective and comprehensive approach to addressing the physiological needs of men and women planning
5 to conceive children.

SUMMARY OF THE INVENTION

The present invention overcomes the deficiencies of currently available approaches to
10 treating male and female infertility. Further, the present invention overcomes the deficiencies of current multi-vitamin and mineral supplements which do not address the specific needs of males and females planning to conceive a child. The
15 present invention overcomes these deficiencies in a safe and effective manner calculated to augment conception, reduce risk of birth defects and generally support the nutritional requirements of developing fetuses and new born infant, as well as
20 contribute to the general health of the mother and father.

The compositions of the invention include critical nutritional components in dosage levels which optimize possibility of conception and fetal
25 development. The compositions are intended for

administration during the period commencing prior to at least two weeks before conception.

Specifically, the present invention provides a nutritional composition for administration to
5 an animal, (e.g., human, mammal or any other animal) during the period commencing prior to at least two weeks before conception, to augment the possibility of conception while enhancing nutritional stores for a developing embryo or
10 fetus prior to and during pregnancy. In one embodiment, the nutritional composition of the present invention comprises about 20 mg to 125 mg per 55 kg of body weight of a vitamin B₆ compound or derivative thereof; about 0.1 mg to 3 mg per
15 55 kg of body weight of a folic acid compound or derivative thereof; and a magnesium compound or derivative thereof in an amount ranging from about 25 mg to 400 mg per 55 kg of body weight. The weight ratio of the folic acid compound or
20 derivative thereof to the vitamin B₆ compound or derivative thereof is about 0.0024-0.1200:1; and the weight ratio of the magnesium compound or derivative thereof to the vitamin B₆ compound or derivative thereof is about 0.2-20:1.

25 An alternative embodiment of the present

invention is a nutritional composition comprising about 20 mg to 125 mg per 55 kg of body weight of a vitamin B₆ compound or derivative thereof; and about 0.1 mg to 3 mg per 55 kg of body weight of a folic acid compound or derivative thereof. The weight ratio of the folic acid compound or derivative thereof to the vitamin B₆ compound or derivative thereof is about 0.0024-0.1200:1.

Another embodiment of the present invention is a nutritional composition comprising about 20 mg to 125 mg per 55 kg of body weight of a vitamin B₆ compound or derivative thereof; and about 100 mg to 1,000 mg per 55 kg of body weight of a calcium compound or derivative thereof. The weight ratio of the calcium compound or derivative thereof to said vitamin B₆ compound or derivative thereof is about 0.25-1:1.

A further embodiment of the present invention is a nutritional composition comprising about 20 mg to 125 mg per 55 kg of body weight of a vitamin B₆ compound or derivative thereof; about 0.1 mg to 3 mg per 55 kg of body weight of a folic acid compound or derivative thereof; and about 100 mg to 1,000 mg per 55 kg of body weight of a calcium compound or derivative thereof per 55 kg of body

weight. The weight ratio of said folic acid compound or derivative thereof to said vitamin B₆ compound or derivative thereof is about 0.0024-0.1200:1; and the weight ratio of said calcium
5 compound or derivative thereof to said vitamin B₆ compound or derivative thereof is about 0.25-1:1.

An even further embodiment of the invention is a nutritional composition for administration to an animal during a period commencing prior to at
10 least two weeks before conception, which comprises: about 0.1 mg to 3 mg per 55 kg of body weight of a folic acid compound or derivative thereof; about 100 mg to 1,000 mg per 55 kg of body weight of a calcium compound or derivative
15 thereof per 55 kg of body weight; and a magnesium compound or derivative thereof in an amount ranging from about 25 mg to 400 mg per 55 kg of body weight. The weight ratio of said folic acid compound or derivative thereof to said calcium
20 compound or derivative thereof is about 0.0001-0.0300:1.

Another embodiment of the present invention includes a nutritional composition comprising about 20 mg to 125 mg per 55 kg of body weight of
25 a vitamin B₆ compound or derivative; about 0.1 mg

to 3 mg per 55 kg of body weight of a folic acid compound or derivative thereof; and about 10 mg to 200 mg per 55 kg of body weight of a fatty acid compound selected from the group consisting of a
5 linoleic acid compound, a linolenic acid compound, and derivatives and mixtures thereof. The weight ratio of said folic acid compound or derivative thereof to said vitamin B₆ compound or derivative thereof is about 0.0024-0.1200:1. The weight
10 ratio of said fatty acid compound to said vitamin B₆ compound or derivative thereof is about 0.08-3.75:1.

Yet another embodiment of the invention provides a nutritional composition comprising
15 about 20 mg to 125 mg per 55 kg of body weight of a vitamin B₆ compound or derivative thereof; about 0.1 mg to 3 mg per 55 kg of body weight of a folic acid compound or derivative thereof; and about 10 mg to 500 mg per 55 kg of body weight of a fatty
20 acid compound selected from the group consisting of a docosahexaenoic acid compound, an arachidonic acid compound, and derivatives and combinations thereof. The weight ratio of said folic acid compound or derivative thereof to said vitamin B₆
25 compound or derivative thereof is about 0.0024-

0.1200:1; and wherein the weight ratio of said fatty acid compound to said vitamin B₆ compound or derivative thereof is about 0.08-25:1.

5 A further embodiment provides a nutritional composition for administration to an animal during a period commencing prior to at least two weeks before conception to augment the possibility of conception while enhancing nutritional stores for a developing embryo or fetus prior to and during
10 pregnancy. The composition comprises about 0.1 mg to 3 mg per 55 kg of body weight of a folic acid compound or derivative thereof; a magnesium compound or derivative thereof in an amount ranging from about 25 mg to 400 mg per 55 kg of
15 body weight; and a vitamin C compound or derivative thereof in an amount ranging from about 25 mg to 600 mg per 55 kg of body weight.

Yet another further embodiment of the invention provides a nutritional composition,
20 which comprises: about 0.1 mg to 3 mg per 55 kg of body weight of a folic acid compound or derivative thereof; a magnesium compound or derivative thereof in an amount ranging from about 25 mg to 400 mg per 55 kg of body weight; and a
25 vitamin E compound or derivative thereof in an

amount ranging from about 10 I.U. to 400 I.U. per 55 kg of body weight.

The present invention also includes methods for augment the possibility of conception while enhancing nutritional stores for a developing embryo or fetus prior to and during pregnancy. In one embodiment, the methods of the present invention include administering to an animal during a period commencing prior to at least two weeks before conception a composition comprising about 20 mg to 125 mg per 55 kg of body weight of a vitamin B₆ compound or derivative thereof; about 0.1 mg to 3 mg per 55 kg of body weight of a folic acid compound or derivative thereof. The weight ratio of said folic acid compound or derivative thereof to said vitamin B₆ compound or derivative thereof is about 0.0024-0.1200:1.

In an alternative embodiment of the invention, a method is provided for increasing the possibility of conception while enhancing nutritional stores for a developing embryo or fetus prior to and during pregnancy, which comprises administering to an animal during a period commencing prior to at least two weeks before conception a composition comprising: about

20 mg to 125 mg per 55 kg of body weight of a vitamin B₆ compound or derivative thereof; and about 100 mg to 1,000 mg per 55 kg of body weight of a calcium compound or derivative thereof. The weight ratio of said calcium compound or derivative thereof to said vitamin B₆ compound or derivative thereof is about 0.25-1:1.

In another embodiment, a method is provided, which comprises administering to an animal during a period commencing prior to at least two weeks before conception a composition including about 20 mg to 125 mg per 55 kg of body weight of a vitamin B₆ compound or derivative thereof; about 0.1 mg to 3 mg per 55 kg of body weight of a folic acid compound or derivative thereof; about 100 mg to 1,000 mg per 55 kg of body weight of a calcium compound or derivative thereof. The weight ratio of said folic acid compound or derivative thereof to said vitamin B₆ compound or derivative thereof is about 0.024-0.1200:1; and the weight ratio of said calcium compound or derivative thereof to said vitamin B₆ compound or derivative thereof is about 0.25-1:1.

In a further embodiment of the invention, a method is provided, which comprises administering

to a male animal and a female animal during a period commencing prior to at least two weeks before conception a composition comprising a nutritional agent selected from the group consisting of a vitamin B₆ compound, a folic acid compound, a magnesium compound, a vitamin C compound, a vitamin E compound, a derivative thereof and a mixture thereof; wherein said male and said female animal are attempting to conceive a child together.

In a still further embodiment of the invention, a method is provided, which comprises: administering to an animal during a period commencing prior to at least two weeks before conception a composition comprising: about 0.1 mg to 3 mg per 55 kg of body weight of a folic acid compound or derivative thereof; and about 100 mg to 1,000 mg per 55 kg of body weight of a calcium compound or derivative thereof.

Thus, the inventive subject matter addresses the specific needs of men and women attempting to conceive through novel compositions and methods. Additional features and advantages of the present invention are described in, and will be apparent from, the detailed description of the presently

preferred embodiments.

DETAILED DESCRIPTION OF THE INVENTION

As used herein, "infertility" refers to the
5 difficulty or inability to conceive during the
course of normal sexual activity, when one year of
unprotected intercourse has elapsed without a
resulting conception.

"Animal" refers to a human, mammal or any
10 other animal.

"Conception" refers to the beginning of
pregnancy as marked by the formation of a zygote.

"Possibility of conception" refers to the
likelihood of conception occurring during normal
15 sexual activity.

"Nutritional stores" refers to the levels of
vitamins, minerals and other nutrients which will
be available for use by the father, mother,
developing embryo, fetus and newborn infant.

20 "Nutritional status" refers to the presence
or absence of any vitamin or mineral deficiency,
or in other words, the extent to which
physiological vitamin and mineral demands are
being satisfied such that deficiency is avoided.

25

The present invention is based, in part, upon the discovery that nutritional requirements vary throughout an individual's lifetime and as a result of various external and internal factors.

5 In particular, certain factors heighten the physiological demand for certain vitamins, minerals and other nutrients and components. Moreover, in some circumstances certain nutritional agents have positive benefits beyond

10 their usual function of maintaining health.

The present inventive subject matter recognizes that there are substantial physiological benefits attained by specifically formulating multi-vitamin and mineral supplements

15 for use by males and females planning to conceive children. Further, the nutritional needs of persons planning to conceive children are different from the nutritional needs of males and females who are not planning to conceive. The

20 products of the invention provide optimum nutritional components and amounts which have been found to alleviate nutritional causes of infertility and provide for optimal health prior to conception.

25 Without being limited by theory, the

compositions and methods of the present invention may be effective because they prevent deficiencies of vitamins, minerals and other nutrients which are necessary to conception. Alternatively, the
5 compositions and methods may be effective because they initiate, stimulate or act as catalysts to reactions having a positive impact on the processes of conception and fetal development.

The multi-vitamin and mineral supplements of
10 the present invention contain specific concentrations of vitamin and minerals for administration to males and females prior to conception to alleviate vitamin and mineral deficiencies which may cause infertility. The
15 present invention also satisfies specific vitamin and mineral requirements, the absence of which have been found to cause birth defects, as well as to provide for general health after conception and during the resultant pregnancy. The formulations
20 of the invention optimize the nutritional benefits of supplementation as required by the physiological stresses of conception.

The nutritional compositions of the present invention are formulated for administration to
25 humans and other animals during the period prior

to and including conception. The effectiveness of the compositions appears to increase in relationship to the length of time between initiation of use and time of conception. Preferably, the compositions are administered during the period commencing prior to at least two weeks before conception. More preferably, the compositions are administered during the period commencing prior to at least four weeks before conception. Even more preferably, the compositions are administered during the period commencing prior to at least twelve weeks prior to conception. Most preferably, the compositions are administered during the period of time commencing prior to at least six months prior to conception.

The nutritional compositions are formulated to augment the possibility of conception. The extent to which the possibility of conception is increased by use of the formulas may be influenced by numerous external factors, such as the following non-limiting examples: stress, alcohol consumption, drug use, poor compliance, and the like. Moreover, the effectiveness of the compositions may vary from individual to

individual, and from couple to couple, for an wide array of reasons, such as genetic predisposition, health factors, and the like, without limitation.

While it is difficult to quantify the likelihood of conception, the average healthy couple may be able to augment conception through use of the present formulations. Moreover, even for couples that could not be classified as average healthy couples, the possibility of conception may be augmented, particularly where the formulations directly impact upon the factor causing the abnormality.

The formulations of the present invention can contain vitamin B₆ or derivatives thereof. Derivatives of vitamin B₆ include compounds formed from vitamin B₆ which are structurally distinct from vitamin B₆, but which retain the active function of vitamin B₆. Such derivatives include, without limitation, salts of vitamin B₆, chelates of vitamin B₆, combinations thereof and the like. The vitamin B₆ may be present in a single form or in various different forms in combination within the present compositions. The specific amount of vitamin B₆ in the compositions is adjusted based on the type of dosage form utilized (i.e.,

immediate release vs. controlled release). In the case of the immediate release compositions, the amounts of vitamin B₆ in the compositions preferably range from about 10 mg to about 75 mg per 55 kg of body weight. More preferably, the amounts of vitamin B₆ in the immediate release compositions range from about 15 mg to about 50 mg per 55 kg of body weight. Even more preferably, the amounts of vitamin B₆ in the immediate release compositions range from about 17 mg to about 25 mg per 55 kg of body weight. Most preferably, the amounts of vitamin B₆ in the immediate release compositions range from about 19 mg to about 21 mg.

The amount of vitamin B₆ present in the controlled release compositions of the present invention, preferably range from about 20 mg to about 150 mg per 55 kg of body weight. More preferably, the amounts of vitamin B₆ present in the controlled release compositions range from about 70 mg to about 125 mg.

The compositions of the present invention may include a folic acid compound or derivative thereof. The derivatives of folic acid include compounds formed from folic acid which are

structurally distinct from folic acid, but which retain the active function of folic acid. Non-limiting examples of such derivatives include salts of folic acid, chelates of folic acid, combinations thereof and the like. Preferably, the amounts of folic acid in the immediate release compositions of the invention range from about 0.1 mg to 3 mg per 55 kg of body weight of a folic acid compound or derivative thereof. More preferably, the amounts of folic acid in the invention range from about 0.5 to about 1 mg per 55 kg of body weight. The amounts of folic acid in the controlled release compositions of the invention preferably range from about 2 mg to about 4 mg per 55 kg of body weight. More preferably, the amounts of folic acid in the controlled release compositions range from about 2.5 to about 3.0 mg per 55 kg of body weight.

A magnesium compound or derivative thereof may be incorporated into the compositions of the present invention. Preferably, the amounts of magnesium in the compositions range from about 10 mg to about 500 mg per 55 kg of body weight. More preferably, the magnesium is in an immediate release form and is present in amounts ranging

from about 15 mg to about 125 mg per 55 kg of body weight. Even more preferably, the magnesium is present in a controlled release form in amounts ranging from about 350 mg to about 450 mg per 55 kg of body weight.

The compositions of the present invention may optionally include a vitamin C compound or derivative thereof. The vitamin C is preferably present in the compositions in amounts ranging from about 5 mg to about 2000 mg per 55 kg of body weight. More preferably, the vitamin C is present in the compositions in amounts ranging from about 60 mg to about 1000 mg per 55 kg of body weight. Even more preferably, the vitamin C is present in a controlled release form in amounts ranging from about 300 mg to about 600 mg per 55 kg of body weight.

The present invention may also additionally include a vitamin E compound or derivative thereof. The vitamin E compound or derivative thereof is preferably present in the compositions in an amount ranging from about 10 I.U. to about 600 I.U. per 55 kg of body weight. More preferably, the vitamin E compound or derivative thereof is present in the compositions in an

immediate release form in an amount ranging from about 5 mg to about 50 mg per 55 kg of body weight. Even more preferably, the vitamin E compound or derivative thereof is present in the compositions in an immediate release form in an amount ranging from about 10 mg to about 30 mg per 55 kg of body weight. Most preferably, the vitamin E compound or derivative thereof is present in the compositions in a controlled release form in an amount ranging from 350 mg to 450 mg per 55 kg of body weight.

The compositions of the present may contain a calcium compound or derivative thereof. The calcium is preferably present in amounts of about 50 mg to about 1,500 mg per 55 kg of body weight. More preferably, the calcium is present in an immediate release form in amounts of about 75 mg to about 1200 mg per 55 kg of body weight. Even more preferably, the calcium is present in an immediate release form in amounts ranging from about 150 mg to about 1000 mg per 55 kg of body weight. Still more preferably the calcium is in a chewable form in amounts ranging from about 75 mg to about 500 mg per 55 kg of body weight.

The compositions of the present invention

optionally contain linoleic acid, linolenic acid, or derivatives or mixtures thereof. Preferably, the linoleic acid, linolenic acid or combination thereof is present in amounts of about 5 mg to 250
5 mg per 55 kg of body weight. More preferably, the linoleic acid, linolenic acid or combination thereof is present in an immediate release form in an amount of ranging from about 5 mg to about 20 mg per 55 kg of body weight. Even more
10 preferably, the linoleic acid, linolenic acid or combination thereof is present in a controlled release form in an amount ranging from about 150 mg to 250 mg per 55 kg of body weight.

The compositions of the present invention may
15 further optionally include docosahexaenoic acid, arachidonic acid or about 10 mg to 200 mg per 55 kg of body weight of a fatty acid compound selected from the group consisting of a docosahexaenoic acid compound, a arachidonic acid
20 compound, and derivatives and mixtures thereof.

Preferred embodiments of the present invention may additionally include a fertility agent, including without limitation, chorionic gonadotropin, clomiphene, gonadorelin, and
25 menotropins.

Particularly, preferred embodiments of the present invention may also further contain one or more ovulatory agents, including without limitation, chorionic gonadotropin, clomiphene, gonadorelin, recombinant human luteinizing hormone, menotropins, progesterone, urofollitropin and combinations thereof.

The weight ratios of various components in the invention are calculated to provide optimal formulations for achieving the objectives of the invention. The weight ratio of the magnesium compound or derivative thereof to said vitamin B₆ compound or derivative thereof in the present invention is preferably about 0.2-20:1. More preferably, the weight ratio of said magnesium compound or derivative thereof to said vitamin B₆ compound or derivative thereof is about 0.5-10:1. Even more preferably, the weight ratio of the magnesium compound or derivative thereof to the vitamin B₆ compound or derivative thereof is about 0.9-5:1.

The weight ratio of folic acid to B₆ in the present compositions is preferably about 0.0024-0.1200:1. More preferably, the weight ratio of folic acid to B₆ in the present compositions is

about 0.0010-0.1100:1. Even more preferably, the weight ratio of folic acid to B₆ in the present compositions is about 0.05-0.09:1. Most preferably, the weight ratio of folic acid to B₆ is 0.02-0.08:1.

The weight ratio of the calcium compound or derivative thereof to said vitamin B₆ compound or derivative thereof in the present invention is preferably about 0.25-1:1. More preferably, the weight ratio of calcium to vitamin B₆ is 0.40-0.90:1. Even more preferably, the weight ratio of calcium to vitamin B₆ is 0.06-0.8:1.

The weight ratio of the linoleic acid, linolenic acid, or derivatives thereof or mixtures thereof to the vitamin B₆ compound or derivative thereof in the present invention is preferably about 0.08-3.75:1. More preferably, the weight ratio of the linoleic acid, linolenic acid, or derivatives thereof or mixtures thereof to the vitamin B₆ compound or derivative thereof in the present invention is 0.15-2.50:1. More preferably, the weight ratio of the linoleic acid, linolenic acid, or derivatives thereof or mixtures thereof to the vitamin B₆ compound or derivative thereof in the present invention is 0.1-1.0:1.

Even more preferably, the weight ratio of the linoleic acid, linolenic acid, or derivatives thereof or mixtures thereof to the vitamin B₆ compound or derivative thereof in the present invention is 0.5-0.9:1.

The weight ratio of the docosahexaenoic acid compound, arachidonic acid compound, or derivatives thereof or mixtures thereof to the vitamin B₆ compound or derivative thereof in the present invention is preferably about 0.08-25:1. More preferably, the weight ratio of the docosohexaenoic acid compound, arachidonic acid compound, or derivatives thereof or mixtures thereof to the vitamin B₆ compound or derivative thereof in the present invention is 1-10:1. Even more preferably, the weight ratio of the docosahexaenoic acid compound, arachidonic acid compound, or derivatives thereof or mixtures thereof to the vitamin B₆ compound or derivative thereof in the present invention is 2-8:1. Most preferably, the weight ratio of the docosahexaenoic acid compound, arachidonic acid compound, or derivatives thereof or mixtures thereof to the vitamin B₆ compound or derivative thereof in the present invention is 4-6:1.

Magnesium compounds which may be incorporated into the present invention include, but are not limited to, magnesium stearate, magnesium carbonate, magnesium oxide, magnesium hydroxide, and magnesium sulfate.

Calcium compounds which may be incorporated into the present invention include, but are not limited to, any of the well known calcium supplements, such as calcium carbonate, calcium sulfate, calcium oxide, calcium hydroxide, calcium apatite, calcium citrate-malate, bone meal, oyster shell, calcium gluconate, calcium lactate, calcium phosphate, calcium levulinate, and the like.

The fatty acids of the present invention may be from any source, including, without limitation, seed oils, fish oil, canola oil, vegetable oil, safflower oil, sunflower oil, olive oil, soybean oil, corn oil, peanut oil, cottonseed oil, chicken fat, lard, palm oil, beef tallow, butter, palm kernel oil, coconut oil, flaxseed oil and evening primrose oil. Non-limiting exemplary fish oil sources include tuna oil, mackerel oil and salmon oil.

It is also possible in the nutritional composition of the present invention for the

dosage form to combine various forms of release, which include, without limitation, immediate release, extended release, pulse release, variable release, controlled release, timed release, 5 sustained release, delayed release, long acting, and combinations thereof. The ability to obtain immediate release, extended release, pulse release, variable release, controlled release, timed release, sustained release, delayed release, 10 long acting characteristics and combinations thereof is performed using well known procedures and techniques available to the ordinary artisan. Each of these specific techniques or procedures for obtaining the release characteristics does not 15 constitute an inventive aspect of this invention.

Any pharmaceutically-acceptable dosage form, and combinations thereof, are contemplated by the invention. Examples of such dosage forms include, without limitation, a chewable tablet, a quick 20 dissolve tablet, an effervescent tablet, reconstitutable powder, elixir, liquid, solution, suspension, emulsion, tablet, multi-layer tablet, bi-layer tablet, capsule, soft gelatin capsule, hard gelatin capsule, caplet, lozenge, chewable 25 lozenge, bead, powder, granules, dispersible

granules, cachets, douche, suppository, cream,
topical, inhalant, aerosol inhalant, patch,
particle inhalant, implant, depot implant,
ingestible, injectable, infusion, a health bar, a
5 liquid, a food and combinations thereof. The
preparation of any of the above dosage forms is
well known in the art.

The following represent examples, without
limitation, of acceptable methods of preparing
10 some of the above-listed dosage forms. For
example, animal feed may be by methods well known
to persons of ordinary skill in the art. Animal
feeds may be prepared by mixing the formulation
with binding ingredients to form a plastic mass.
15 The mass is then extruded under high pressure to
form tubular (or "spaghetti-like") structures that
are cut to pellet size and dried.

Quick dissolve tablets may be prepared, for
example, without limitation, by mixing the
20 formulation with agents such as sugars and
cellulose derivatives, which promote dissolution
or disintegration of the resultant tablet after
oral administration, usually within 30 seconds.

Cereal coatings may be prepared, for example,
25 without limitation, by passing the cereal

formulation, after it has been formed into pellets, flakes, or other geometric shapes, under a precision spray coating device to deposit a film of active ingredients, plus excipients onto the
5 surface of the formed elements. The units thus treated are then dried to form a cereal coating.

For example, health bars may be prepared, without limitation, by mixing the formulation plus excipients (e.g., binders, fillers, flavors,
10 colors, etc.) to a plastic mass consistency. The mass is then either extended or molded to form "candy bar" shapes that are then dried or allowed to solidify to form the final product.

Soft gel or soft gelatin capsules may be prepared, for example, without limitation, by dispersing the formulation in an appropriate vehicle (vegetable oils are commonly used) to form a high viscosity mixture. This mixture is then encapsulated with a gelatin based film using
15 technology and machinery known to those in the soft gel industry. The industrial units so formed are then dried to constant weight.

Chewable tablets, for example, without limitation, may be prepared by mixing the formulations with excipients designed to form a
25

relatively soft, flavored, tablet dosage form that is intended to be chewed rather than swallowed. Conventional tablet machinery and procedures, that is both direct compression and granulation, i.e.,
5 or slugging, before compression, can be utilized. Those individuals involved in pharmaceutical solid dosage form production are well versed in the processes and the machinery used as the chewable dosage form is a very common dosage form in the
10 pharmaceutical industry.

Film coated tablets, for example, without limitation, may be prepared by coating tablets using techniques such as rotating pan coating methods or air suspension methods to deposit a
15 contiguous film layer on a tablet. This procedure is often done to improve the aesthetic appearance of tablets, but may also be done to improve the swallowing of tablets, or to mask an obnoxious odor or taste, or to improve the usual properties
20 of an unsightly uncoated tablet.

Compressed tablets, for example, without limitation, may be prepared by mixing the formulation with excipients intended to add binding qualities to disintegration qualities.
25 The mixture is either directly compressed or

granulated then compressed using methods and machinery quite well known to those in the industry. The resultant compressed tablet dosage units are then packaged according to market need,
5 i.e., unit dose, rolls, bulk bottles, blister packs, etc.

The present invention contemplates nutritional compositions formulated for administration by any route, including without
10 limitation, oral, buccal, sublingual, rectal, parenteral, topical, inhalational, injectable and transdermal. The physicochemical properties of nutritional compositions, their formulations, and the routes of administration are important in
15 absorption. Absorption refers to the process of nutritional composition movement from the site of administration toward the systemic circulation. Most orally administered nutritional compositions are in the form of tablets or capsules primarily
20 for convenience, economy, stability, and patient acceptance. They must disintegrate and dissolve before absorption can occur. Using the present invention with any of the above routes of administration or dosage forms is performed using
25 well known procedures and techniques available to

the ordinary skilled artisan.

The present invention contemplates the use of pharmaceutically acceptable carriers which may be prepared from a wide range of materials. Without
5 being limited thereto, such materials include diluents, binders and adhesives, lubricants, plasticizers, disintegrants, colorants, bulking substances, flavorings, sweeteners and miscellaneous materials such as buffers and
10 adsorbents in order to prepare a particular medicated composition.

Binders may be selected from a wide range of materials such as hydroxypropylmethylcellulose, ethylcellulose, or other suitable cellulose
15 derivatives, povidone, acrylic and methacrylic acid co-polymers, pharmaceutical glaze, gums, milk derivatives, such as whey, starches, and derivatives, as well as other conventional binders well known to persons skilled in the art.
20 Exemplary non-limiting solvents are water, ethanol, isopropyl alcohol, methylene chloride or mixtures and combinations thereof. Exemplary non-limiting bulking substances include sugar, lactose, gelatin, starch, and silicon dioxide.

25 The plasticizers used in the dissolution

modifying system are preferably previously dissolved in an organic solvent and added in solution form. Preferred plasticizers may be selected from the group consisting of diethyl
5 phthalate, diethyl sebacate, triethyl citrate, cronotic acid, propylene glycol, butyl phthalate, dibutyl sebacate, castor oil and mixtures thereof, without limitation. As is evident, the plasticizers may be hydrophobic as well as
10 hydrophilic in nature. Water-insoluble hydrophobic substances, such as diethyl phthalate, diethyl sebacate and castor oil are used to delay the release of water-soluble vitamins, such as vitamin B₆ and vitamin C. In contrast,
15 hydrophilic plasticizers are used when water-insoluble vitamins are employed which aid in dissolving the encapsulated film, making channels in the surface, which aid in nutritional composition release.

20 The dosage forms of the present invention may involve the administration of a nutritional composition in a partial, i.e., fractional dose, one or more times during a 24 hour period, a single dose during a 24 hour period of time, a
25 double dose during a 24 hour period of time, or

more than a double dose during a 24 hour period of time. Fractional, double or multiple doses may be taken simultaneously or at different times during the 24 hour period.

5 The compositions of the present invention are intended for use by humans and other animals, and both males and females. The dosages are adjusted according to body weight and thus are set forth herein on a per body weight basis. For example,
10 if the formula specifies a range of 20-125 mg for a 55 kg individual, that range would be adjusted for a 35 kg individual to 13-80 mg (e.g., the lower range limit = $(35 \text{ kg} / 55 \text{ kg}) * 20 \text{ mg} = 12.6 \text{ mg}$, or about 13 mg). Decimal amount may be
15 rounded to the nearest whole number. In the above manner, the present compositions may be thus adapted to be suitable for any individual, including any animal, regardless of size.

 Moreover, the formulations can be further
20 adapted based upon the specific needs, genetic predispositions or identified deficiencies of the individual planning to conceive. Moreover, the present compositions can be used as one component of a prescribed therapy. Preferably, the
25 compositions are used by both the male and female

planning to conceive a child together.

Pharmaceutically acceptable calcium compounds include, but are not limited to, any of the well known calcium supplements, such as calcium carbonate, calcium sulfate, calcium oxide, calcium hydroxide, calcium apatite, calcium citrate-malate, bone meal, oyster shell, calcium gluconate, calcium lactate, calcium phosphate, calcium levulinate, and the like.

10 The present invention includes methods for increasing the possibility of conception while improving the nutritional stores of the mother prior to conception and enhancing nutritional stores for a developing embryo or fetus prior to and during pregnancy. The methods of the invention comprise administering to a male or a female during a period commencing prior to at least two weeks before conception a composition comprising a vitamin B₆ compound, a folic acid compound, a magnesium compound, a calcium compound, a vitamin C compound, a vitamin E compound, a derivative thereof and a mixture thereof. Preferably, the methods of the invention comprise administering the above described composition to both a male and a female planning

15

20

25

to conceive a child together. More preferably, the methods of the invention comprise administering any of the above described compositions to males and/or females.

5 In one preferred embodiment of the invention, the compositions of the invention are provided to a male or female in a blister pack. In a particularly preferred embodiment of the invention, the multi-vitamin and mineral
10 supplements described above are provided to both a male and female in a blister pack with indicia identifying which supplement is for the male and which supplement is for the female, where the male and female are planning to conceive a child
15 together. All of the compositions of the present invention may further include or be accompanied by indicia allowing men and women to identify the compositions as products for persons planning to conceive children.

20 The methods of the invention may involve the administration of a nutritional composition in a single dose during a 24 hour period of time, a double dose during a 24 hour period of time, or more than two dose during a 24 hour period of
25 time. The dose may be taken simultaneously or at

different times depending upon the prescribed dose.

The foregoing is considered as illustrative only of the principles of the invention. Further, since numerous modifications and changes will readily occur to those skilled in the art, it is not desired to limit the invention to the exact construction and operation shown and described, and accordingly all suitable modifications and equivalents may be resorted to, falling within the scope of the invention. The following examples are illustrative of preferred embodiments of the invention and are not to be construed as limiting the invention thereto.

EXAMPLES

Preparation of Nutritional Conception Compositions

Example 1

The following compositions are used to prepare nutritional conception products for administration to men and women during a period commencing prior to at least two weeks before conception:

Table I

COMPONENT	FORMULA I	CHEWABL E FORMULA	CONTROLLE D RELEASE
Vitamin E, I.U.	10-100	10-100	100-400
Vitamin C, mg	25-100	25-100	100-2,000
Vitamin B ₆ , mg	20-75	20-75	75-125
Folic Acid, mg	0.1-1.5	0.1-1.5	1.5-3
Calcium, mg	100-300	100-300	300-1,500
Magnesium, mg	25-100	25-100	100-400
Linolenic Acid, mg	10-100	-	100-200
Linoleic Acid, mg	10-100	-	100-200
Docosahexaenoic Acid, mg	10-500	-	10-500

Tablets incorporating the above formulations are prepared using conventional methods and materials known in the pharmaceutical art. The resulting nutritional conception composition tablets were recovered and stored for future use.

Example II

The following compositions are used to prepare nutritional conception products for administration during a period commencing prior to at least two weeks before conception:

Table II

	Component (in mg unless otherwise indicated)	Formula I	Chewable Formula	Controlled Release
5	Beta Carotene I.U.	3,000	5,000	3,000
	Vitamin E, I.U.	30	10	400
	Vitamin C,	60	25	600
10	Vitamin B ₁	3.0	3	3
	Vitamin B ₂	3.4	3.4	3.4
	Vitamin B ₃	20	20	20
	Vitamin B ₆	20	20	125
	Vitamin B ₁₂ , mcg	12	12	12
15	Folic Acid	1	1	3
	Calcium	200	100	1,000
	Elemental Iron	30	30	30
	Copper	2	2	2
	Zinc	15	15	15
20	Magnesium	100	25	400
	Microcrystalline Cellulose	180	180	-
	Croscarmellose Sodium	15	-	-
25	Stearic Acid	65	-	-
	Mg Stearate	9	9	10
	Linolenic Acid	10	-	200
	Linoleic Acid	10	-	200
	Docosahexaenoic Acid	500	-	10
30	Ethyl Cellulose	-	-	150

Tablets incorporating the above formulations are prepared using conventional methods and materials known in the pharmaceutical art. The resulting nutritional conception composition tablets are recovered and stored for future use.

Example III

The following compositions are used to prepare nutritional conception products for administration during a period commencing prior to at least two weeks before conception:

Table III

COMPONENT	PER 35 KG OF BODY WEIGHT	PER 55 KG OF BODY WEIGHT	PER 75 KG OF BODY WEIGHT
Vitamin E, I.U.	6-255	10-400	14-546
Vitamin C, mg	16-382	25-600	34-818
Vitamin B ₆ , mg	13-80	20-125	27-171
Folic Acid, mg	0.06-1.9	0.1-3.0	0.14-4.1
Calcium, mg	64-636	100- 1,000	136-1,364
Magnesium, mg	16-255	25-400	34-546
Linolenic Acid, mg	6-127	10-200	14-273
Linoleic Acid, mg	6-127	10-200	14-273
Docosahexaenoic Acid, mg	6-318	10-500	14-682

Tablets incorporating the above formulations are prepared using conventional methods and materials known in the pharmaceutical art. The resulting nutritional conception composition tablets are recovered and stored for future use.

The invention being thus described, it will be apparent that the same may be varied in many ways. Such variations are not to be regarded as a departure from the spirit and scope of the invention, and all such modifications are intended to be within the scope of the appended claims.

15

We Claim:

1. A nutritional composition for administration to an animal during a period commencing prior to at least two weeks before
5 conception to augment the possibility of conception while enhancing nutritional stores for a developing embryo or fetus prior to and during pregnancy, which comprises:

about 20 mg to 125 mg per 55 kg of said
10 animal's body weight of a vitamin B₆ compound or derivative thereof;

about 0.1 mg to 3 mg per 55 kg of said animal's body weight of a folic acid compound or derivative thereof;

15 a magnesium compound or derivative thereof in an amount ranging from about 25 mg to 400 mg per 55 kg of said animal's body weight;

wherein the weight ratio of said folic acid compound or derivative thereof to said vitamin B₆
20 compound or derivative thereof is about 0.0024-0.1200:1; and

wherein the weight ratio of said magnesium compound or derivative thereof to said vitamin B₆ compound or derivative thereof is about 0.2-20:1.

25

2. The nutritional composition of claim 1,
wherein said animal is of a female gender.

5 3. The nutritional composition of claim 1,
wherein said nutritional composition is in an oral
dosage form.

10 4. The nutritional composition of claim 3,
wherein said oral dosage form is selected from the
group consisting of immediate release, extended
release, pulsed release, delayed release, timed
release, variable release, controlled release and
combinations thereof.

15 5. The nutritional composition of claim 3,
wherein said oral dosage form is selected from the
group consisting of a chewable tablet, a quick
dissolve tablet, an effervescent tablet, a hard
20 gelatin capsule, a soft gelatin capsule, a
reconstitutable powder, a suspension, an elixir,
a caplet, a health bar, a liquid, a food and
combinations thereof.

25 6. The nutritional composition of claim 3,

wherein said oral dosage form is a multiple layered tablet.

7. The nutritional composition of claim 1,
5 wherein said nutritional composition is administered once during a twenty four hour period of time.

8. The nutritional composition of claim 1,
10 wherein said nutritional composition is administered at least twice during a twenty four hour period of time.

9. The nutritional composition of claim 1,
15 wherein said nutritional composition further comprises a vitamin C compound or derivative thereof in an amount ranging from about 25 mg to 2000 mg per 55 kg of said animal's body weight.

20 10. The nutritional composition of claim 1, wherein said nutritional composition further comprises a vitamin E compound or derivative thereof in an amount ranging from about 10 I.U. to 400 I.U. per 55 kg of said animal's body weight.

25

11. The nutritional composition of claim 1,
wherein said nutritional composition additionally
contains a fertility agent.

5 12. The nutritional composition of claim 1,
wherein said nutritional composition additionally
contains an ovulatory agent.

10 13. A nutritional composition for
administration to an animal during a period
commencing prior to at least two weeks before
conception, which comprises:

15 about 20 mg to 125 mg per 55 kg of said
animal's body weight of a vitamin B₆ compound or
derivative thereof;

 about 0.1 mg to 3 mg per 55 kg of said
animal's body weight of a folic acid compound or
derivative thereof; and

20 wherein the weight ratio of said folic acid
compound or derivative thereof to said vitamin B₆
compound or derivative thereof is about 0.0024-
0.1200:1.

25 14. The nutritional composition of claim 13,
wherein said animal is of a female gender.

15. The nutritional composition of claim 13,
wherein said animal is of a male gender.

16. The nutritional composition of claim 13,
5 wherein said nutritional composition further
comprises a magnesium compound or derivative
thereof in an amount ranging from about 25 mg to
400 mg per 55 kg of said animal's body weight.

10 17. The nutritional composition of claim 13,
wherein said nutritional composition is
administered once during a twenty four hour period
of time.

15 18. The nutritional composition of claim 13,
wherein said nutritional composition is
administered at least twice during a twenty four
hour period of time.

20 19. The nutritional composition of claim 13,
wherein said composition is administered to both
a male animal and a female animal attempting to
conceive a child together.

25 20. A nutritional composition for

administration to an animal during a period commencing prior to at least two weeks before conception, which comprises:

5 about 20 mg to 125 mg per 55 kg of said animal's body weight of a vitamin B₆ compound or derivative thereof;

 about 100 mg to 1,000 mg per 55 kg of said animal's body weight of a calcium compound or derivative thereof; and

10 wherein the weight ratio of said calcium compound or derivative thereof to said vitamin B₆ compound or derivative thereof is about 0.25-1:1.

21. The nutritional composition of claim 20,
15 wherein said nutritional composition is in an oral dosage form.

22. The nutritional composition of claim 21,
 wherein said oral dosage form is selected from the
20 group consisting of a chewable tablet, a quick dissolve tablet, an effervescent tablet, a hard gelatin capsule, a soft gelatin capsule, a reconstitutable powder, a suspension, an elixir, a caplet, a health bar, a liquid, a food and
25 combinations thereof.

23. The nutritional composition of claim 21,
wherein said oral dosage form is selected from the
group consisting of immediate release, extended
release, pulsed release, delayed release, timed
5 release, variable release, controlled release and
combinations thereof.

24. The nutritional composition of claim 20,
wherein said nutritional composition additionally
10 includes a folic acid compound or derivative
thereof in an amount of about 0.1 mg to 3.0 mg.

25. The nutritional composition of claim 20,
wherein said nutritional composition is
15 administered once during a twenty four hour period
of time.

26. The nutritional composition of claim 20,
wherein said nutritional composition is
20 administered at least twice during a twenty four
hour period of time.

27. The nutritional composition of claim 20,
wherein said nutritional composition further
25 comprises a magnesium compound or derivative

thereof in an amount ranging from about 25 mg to 400 mg per 55 kg of said animal's body weight.

28. The nutritional composition of claim 20,
5 wherein said composition is administered to both a male animal and a female animal attempting to conceive a child together.

29. A nutritional composition for
10 administration to an animal during a period commencing prior to at least two weeks before conception, which comprises:

a) about 20 mg to 125 mg per 55 kg of said
animal's body weight of a vitamin B₆ compound or
15 derivative thereof;

b) about 0.1 mg to 3 mg per 55 kg of said
animal's body weight of a folic acid compound or
derivative thereof;

c) about 100 mg to 1,000 mg per 55 kg of
20 said animal's body weight of a calcium compound or derivative thereof per 55 kg of said animal's body weight;

wherein the weight ratio of said folic acid
compound or derivative thereof to said vitamin B₆
25 compound or derivative thereof is about 0.0024-

0.1200:1; and

wherein the weight ratio of said calcium compound or derivative thereof to said vitamin B₆ compound or derivative thereof is about 0.25-1:1.

5

30. The nutritional composition of claim 29, wherein said composition is in an oral dosage form.

10

31. The nutritional composition of claim 30, wherein said oral dosage form is selected from the group consisting of a chewable tablet, a quick dissolve tablet, an effervescent tablet, a hard gelatin capsule, a soft gelatin capsule, a reconstitutable powder, a suspension, an elixir, a caplet, a health bar, a liquid, a food and combinations thereof.

15

32. The nutritional composition of claim 30, wherein said oral dosage form is selected from the group consisting of immediate release, extended release, pulsed release, delayed release, timed release, variable release, controlled release and combinations thereof.

20

25

33. The nutritional composition of claim 30, wherein said nutritional composition is administered once during a twenty four hour period of time.

5

34. The nutritional composition of claim 30, wherein said nutritional composition is administered at least twice during a twenty four hour period of time.

10

35. A nutritional composition for administration to an animal during a period commencing prior to at least two weeks before conception, which comprises:

15 about 0.1 mg to 3 mg per 55 kg of said animal's body weight of a folic acid compound or derivative thereof;

 about 100 mg to 1,000 mg per 55 kg of said animal's body weight of a calcium compound or derivative thereof per 55 kg of said animal's body weight;

20

 a magnesium compound or derivative thereof in an amount ranging from about 25 mg to 400 mg per 55 kg of said animal's body weight; and

25 wherein the weight ratio of said folic acid

compound or derivative thereof to said calcium compound or derivative thereof is about 0.0001-0.0300:1.

5 36. A nutritional composition for administration to an animal during a period commencing prior to at least two weeks before conception, which comprises:

10 a) about 20 mg to 125 mg per 55 kg of said animal's body weight of a vitamin B₆ compound or derivative;

 b) about 0.1 mg to 3 mg per 55 kg of said animal's body weight of a folic acid compound or derivative thereof;

15 c) about 10 mg to 200 mg per 55 kg of said animal's body weight of a fatty acid compound selected from the group consisting of a linoleic acid compound, a linolenic acid compound, and derivatives and mixtures thereof;

20 wherein the weight ratio of said folic acid compound or derivative thereof to said vitamin B₆ compound or derivative thereof is about 0.0024-0.1200:1; and

25 wherein the weight ratio of said fatty acid compound to said vitamin B₆ compound or derivative

thereof is about 0.08-3.75:1.

37. The nutritional composition of claim 36,
wherein said nutritional composition is in an oral
5 dosage form.

38. The nutritional composition of claim 37,
wherein said oral dosage form is selected from the
group consisting of a chewable tablet, a quick
10 dissolve tablet, an effervescent tablet, a hard
gelatin capsule, a soft gelatin capsule, a
reconstitutable powder, a suspension, an elixir,
a caplet, a health bar, a liquid, a food and
combinations thereof.

15 39. The nutritional composition of claim 37,
wherein said oral dosage form is selected from the
group consisting of immediate release, extended
release, pulsed release, delayed release, timed
20 release, variable release, controlled release and
combinations thereof.

40. The nutritional composition of claim 36,
wherein said nutritional composition is
25 administered once during a twenty four hour period

of time.

41. The nutritional composition of claim 36,
wherein said nutritional composition is
5 administered at least twice during a twenty four
hour period of time.

42. A nutritional composition for
administration to an animal during a period
10 commencing prior to at least two weeks before
conception, which comprises:

a) about 20 mg to 125 mg per 55 kg of said
animal's body weight of a vitamin B₆ compound or
derivative thereof;

15 b) about 0.1 mg to 3 mg per 55 kg of said
animal's body weight of a folic acid compound or
derivative thereof;

c) about 10 mg to 500 mg per 55 kg of said
animal's body weight of a fatty acid compound
20 selected from the group consisting of a
docosahexaenoic acid compound, an arachidonic acid
compound, and derivatives and combinations
thereof;

wherein the weight ratio of said folic acid
25 compound or derivative thereof to said vitamin B₆

compound or derivative thereof is about 0.0024-0.1200:1; and

wherein the weight ratio of said fatty acid compound to said vitamin B₆ compound or derivative thereof is about 0.08-25:1.

43. The nutritional composition of claim 42, wherein said composition is in an oral dosage form.

10

44. The nutritional composition of claim 43, wherein said oral dosage form is selected from the group consisting of a chewable tablet, a quick dissolve tablet, an effervescent tablet, a hard gelatin capsule, a soft gelatin capsule, a reconstitutable powder, a suspension, an elixir, a caplet, a health bar, a liquid, a food and combinations thereof.

15

45. The nutritional composition of claim 43, wherein said oral dosage form is selected from the group consisting of immediate release, extended release, pulsed release, delayed release, timed release, variable release, controlled release and combinations thereof.

20

25

46. The nutritional composition of claim 42, wherein said nutritional composition is administered once during a twenty four hour period of time.

5

47. The nutritional composition of claim 42, wherein said nutritional composition is administered at least twice during a twenty four hour period of time.

10

48. A nutritional composition for administration to an animal during a period commencing prior to at least two weeks before conception to augment the possibility of conception while enhancing nutritional stores for a developing embryo or fetus prior to and during pregnancy, which comprises:

15

about 0.1 mg to 3 mg per 55 kg of said animal's body weight of a folic acid compound or derivative thereof;

20

a magnesium compound or derivative thereof in an amount ranging from about 25 mg to 400 mg per 55 kg of said animal's body weight; and

a vitamin C compound or derivative thereof in an amount ranging from about 25 mg to 600 mg per

25

55 kg of said animal's body weight.

49. The nutritional composition of claim 48,
wherein said animal is of a male gender.

5

50. The nutritional composition of claim 48,
wherein said composition is in an oral dosage
form.

10

51. The nutritional composition of claim 50,
wherein said oral dosage form is selected from the
group consisting of a chewable tablet, a quick
dissolve tablet, an effervescent tablet, a hard
gelatin capsule, a soft gelatin capsule, a
15 reconstitutable powder, a suspension, an elixir,
a caplet, a health bar, a liquid, a food and
combinations thereof.

20

52. The nutritional composition of claim 50,
wherein said oral dosage form is selected from the
group consisting of immediate release, extended
release, pulsed release, delayed release, timed
release, variable release, controlled release and
combinations thereof.

25

53. The nutritional composition of claim 48, wherein said nutritional composition is administered once during a twenty four hour period of time.

5

54. The nutritional composition of claim 48, wherein said nutritional composition is administered at least twice during a twenty four hour period of time.

10

55. A nutritional composition for administration to an animal during a period commencing prior to at least two weeks before conception to augment the possibility of conception while enhancing nutritional stores for a developing embryo or fetus prior to and during pregnancy, which comprises:

15

about 0.1 mg to 3 mg per 55 kg of said animal's body weight of a folic acid compound or derivative thereof;

20

a magnesium compound or derivative thereof in an amount ranging from about 25 mg to 400 mg per 55 kg of said animal's body weight; and

a vitamin E compound or derivative thereof in an amount ranging from about 10 I.U. to 400 I.U.

25

per 55 kg of said animal's body weight.

56. The nutritional composition of claim 55,
wherein said animal is of a male gender.

5

57. The nutritional composition of claim 55,
wherein said composition is in an oral dosage
form.

10

58. The nutritional composition of claim 57,
wherein said oral dosage form is selected from the
group consisting of a chewable tablet, a quick
dissolve tablet, an effervescent tablet, a hard
gelatin capsule, a soft gelatin capsule, a
15 reconstitutable powder, a suspension, an elixir,
a caplet, a health bar, a liquid, a food and
combinations thereof.

15

59. The nutritional composition of claim 57,
20 wherein said oral dosage form is selected from the
group consisting of immediate release, extended
release, pulsed release, delayed release, timed
release, variable release, controlled release and
combinations thereof.

25

60. The nutritional composition of claim 55, wherein said nutritional composition is administered once during a twenty four hour period of time.

5

61. The nutritional composition of claim 55, wherein said nutritional composition is administered at least twice during a twenty four hour period of time.

10

62. A method for increasing the possibility of conception while enhancing nutritional stores for a developing embryo or fetus prior to and during pregnancy, which comprises:

15 administering to an animal during a period commencing prior to at least two weeks before conception a composition comprising:

about 20 mg to 125 mg per 55 kg of said animal's body weight of a vitamin B₆ compound or derivative thereof;

20

about 0.1 mg to 3 mg per 55 kg of said animal's body weight of a folic acid compound or derivative thereof; and

wherein the weight ratio of said folic acid compound or derivative thereof to said vitamin B₆

25

compound or derivative thereof is about 0.0024-
0.1200:1.

63. The method of claim 61, wherein said
5 animal is of a male gender.

64. The method of claim 61, wherein said
animal is of a female gender.

10 65. The method of claim 61, wherein said
nutritional composition is in an oral dosage form.

66. The method of claim 65, wherein said
oral dosage form is selected from the group
15 consisting of a chewable tablet, a quick dissolve
tablet, an effervescent tablet, a hard gelatin
capsule, a soft gelatin capsule, a reconstitutable
powder, a suspension, an elixir, a caplet, a
health bar, a liquid, a food and combinations
20 thereof.

67. The method of claim 65, wherein said
oral dosage form is selected from the group
consisting of immediate release, extended release,
25 pulsed release, delayed release, timed release,

variable release, controlled release and combinations thereof.

5 68. The method of claim 61, wherein said nutritional composition is administered once during a twenty four hour period of time.

10 69. The method of claim 61, wherein said nutritional composition is administered at least twice during a twenty four hour period of time.

15 70. The method of claim 61, wherein said nutritional composition further comprises a vitamin C compound or derivative thereof in an amount ranging from about 25 mg to 400 mg per 55 kg of said animal's body weight.

20 71. The method of claim 61, wherein said nutritional composition further comprises a vitamin E compound or derivative thereof in an amount ranging from about 10 mg to 400 mg per 55 kg of said animal's body weight.

25 72. The method of claim 61, wherein said nutritional composition additionally contains a

fertility agent.

73. The method of claim 61, wherein said
nutritional composition additionally contains an
5 ovulatory agent.

74. The method of claim 61, wherein said
composition is administered about 8 hours prior to
sexual intercourse.

10

75. A method for increasing the possibility
of conception while enhancing nutritional stores
for a developing embryo or fetus prior to and
during pregnancy, which comprises:

15 administering to an animal during a period
commencing prior to at least two weeks before
conception a composition comprising:

about 20 mg to 125 mg per 55 kg of said
animal's body weight of a vitamin B₆ compound or
20 derivative thereof;

about 100 mg to 1,000 mg per 55 kg of said
animal's body weight of a calcium compound or
derivative thereof; and

wherein the weight ratio of said calcium
25 compound or derivative thereof to said vitamin B₆

compound or derivative thereof is about 0.25-1:1.

76. The method of claim 75, wherein said nutritional composition is administered once
5 during a twenty four hour period of time.

77. The method of claim 75, wherein said nutritional composition is administered at least twice during a twenty four hour period of time.

10

78. A method for increasing the possibility of conception while enhancing nutritional stores for a developing embryo or fetus prior to and during pregnancy, which comprises:

15 administering to an animal during a period commencing prior to at least two weeks before conception a composition comprising:

a) about 20 mg to 125 mg per 55 kg of said animal's body weight of a vitamin B₆ compound or
20 derivative thereof;

b) about 0.1 mg to 3 mg per 55 kg of said animal's body weight of a folic acid compound or derivative thereof;

c) about 100 mg to 1,000 mg per 55 kg of
25 said animal's body weight of a calcium compound or

derivative thereof;

wherein the weight ratio of said folic acid compound or derivative thereof to said vitamin B₆ compound or derivative thereof is about 0.024-
5 0.1200:1; and

wherein the weight ratio of said calcium compound or derivative thereof to said vitamin B₆ compound or derivative thereof is about 0.25-1:1.

10 79. The method of claim 78, wherein said nutritional composition is administered once during a twenty four hour period of time.

15 80. The method of claim 78, wherein said nutritional composition is administered at least twice during a twenty four hour period of time.

20 81. A method for increasing the possibility of conception while improving nutritional status for a mother prior to conception and enhancing nutritional stores for a developing embryo or fetus prior to and during pregnancy, which comprises:

25 administering to a male animal and a female animal during a period commencing prior to at

least two weeks before conception a composition comprising a nutritional agent selected from the group consisting of a vitamin B₆ compound, a folic acid compound, a magnesium compound, a vitamin C
5 compound, a vitamin E compound, a derivative thereof and a mixture thereof; and

wherein said male and said female animal are attempting to conceive a child.

10 82. The method of claim 81, wherein said composition is provided to said male animal and said female animal on a blister pack with indicia for both the male animal and female animal.

15 83. A method for increasing the possibility of conception while enhancing nutritional stores for a developing embryo or fetus prior to and during pregnancy, which comprises:

administering to an animal during a period
20 commencing prior to at least two weeks before conception a composition comprising:

about 0.1 mg to 3 mg per 55 kg of said
animal's body weight of a folic acid compound or
derivative thereof; and

25 about 100 mg to 1,000 mg per 55 kg of said

animal's body weight of a calcium compound or derivative thereof.

84. The method of claim 83, wherein said
5 nutritional composition is administered once
during a twenty four hour period of time.

85. The method of claim 83, wherein said
nutritional composition is administered at least
10 twice during a twenty four hour period of time.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US00/11681**A. CLASSIFICATION OF SUBJECT MATTER**

IPC(7) : A61K 33/06, 31/20, 31/22, 31/52, 31/55

US CL : 514/185, 261, 549, 558; 424/682

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 514/185, 261, 549, 558; 424/682

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

MEDLINE, CAPLUS, BIOSIS

search terms: fertility, infertility, nutrition, VITAMIN E, B6, B12, FOLIC ACID,

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US 5,231,085 A (ALEXANDER et al) 27 July 1993 (27.07.93), column 1, lines 14-16 and column 6, lines 1-67.	1-85
Y,P	US 5,922,704 A (BLAND) 13 July 1999 (13.07.99), see entire document, especially columns 6-8.	1-85
Y	Database Medline on STN, US National Library of Medicine, (Bethesda, MD, USA), No. 86054328, VAN DER SPUIY, Z. M. 'Nutrition and Reproduction,' abstract, Clinics in Obstetrics and Gynaecology, September 1985, 12(3), 579-604.	1-85



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
B earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*A* document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means	
P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

09 JUNE 2000

Date of mailing of the international search report

17 AUG 2000

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer:

SHENGJUN WANG

Telephone No. (703) 308-1235

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US00/11681

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	Database CAPLUS on STN, (Columbus. OH, USA), No. 91:173855, OCETKIEWICZ et al., 'Study of the suitability of premixes in rabbit,' abstract, Rocz. Nauk. Zootech. 1977, 4 (2), 161-173, (Polish).	1-85